arXiv:1105.5740v3 [math.PR] 17 Oct 2011

Quenched invariance principle for simple random walk on discrete point processes

Naoki KUBOTA[*]

Abstract

We consider the simple random walk on random graphs generated by discrete point processes. This random graph has a random subset of a cubic lattice as the vertices and lines between any consecutive vertices on lines parallel to each coordinate axis as the edges. Under the assumption that discrete point processes are finitely dependent and stationary, we prove that the quenched invariance principle holds, that is, for almost every configuration of a point process, the path distribution of the walk converges weakly to that of a Brownian motion.

1 Introduction

Random walks in random environment constitute one of the basic models of random motions in random media. On the validity of the quenched invariance principle, the ballistic random walk in random environment and the random walk among random conductances have been intensively investigated in recent years. These models treat random walks with bounded jumps and i.i.d. configurations. Our interest now is the quenched invariance principle for models that random walks have non-bounded jumps and configurations are not independent. We consider a simple random walk on discrete point processes on $\mathbb{Z}^d$, that is, a simple random walk on the random graph whose vertices are a random subset of $\mathbb{Z}^d$ and whose edges are the lines between any consecutive vertices on the lines parallel to each coordinate axis. This model was introduced in [7], and its law of large numbers and central limit theorem are shown here. It is open whether to approve the quenched invariance principle for this model, see [7, Section 11]. The aim of this paper is to prove the quenched invariance principle when point processes are finitely dependent and stationary.

We now describe the setting in more detail. Let $\Omega := \{0,1\}^{\mathbb{Z}^d}$ and denote by $\omega = (\omega(x))_{x\in\mathbb{Z}^d}$ an element of Ω. The space Ω is equipped with the canonical product σ-field $\mathcal{G}$, the canonical shift $T_y\omega(x) := \omega(x+y)$ for $x, y \in \mathbb{Z}^d$ and a probability measure $\mathbb{Q}$. Let $\Omega_x := \{\omega \in \Omega; \omega(x) = 1\}$ for $x \in \mathbb{Z}^d$ and we shall assume throughout the paper that $\mathbb{Q}$ satisfies the following conditions:

(A1) $0 < \mathbb{Q}(\Omega_0) < 1$ holds,

(A2) There is a positive constant ℓ such that if A, $B \subset \mathbb{Z}^d$ satisfy $\inf\{|x-y|; x \in A, y \in B\} \geq \ell$ then $\sigma(\omega(x); x \in A)$ and $\sigma(\omega(x); x \in B)$ are independent,

---

[*] Department of Mathematics, Graduate School of Science and Technology, Nihon University, mail: kubota@grad.math.cst.nihon-u.ac.jp

(A3) $\mathbb{Q}$ is stationary with respect to canonical shifts $(T_x)_{x\in\mathbb{Z}^d}$.

Then we can define the probability measure $\mathbb{P}$ on Ω_0 as follows:

$$\mathbb{P}(A) := \mathbb{Q}(A|\Omega_0), \qquad A \in \mathcal{G},$$

and denote by $E_{\mathbb{Q}}$ and $\mathbb{E}$ the expectation with respect to $\mathbb{Q}$ and $\mathbb{P}$, respectively. Let $\omega \in \Omega$ and set $\mathcal{P}(\omega) := \{x \in \mathbb{Z}^d; \omega(x) = 1\}$. It is immediate to see from the assumptions (A1)-(A3) that $\gamma_e(\omega) := \inf\{k \geq 1; \omega(ke) = 1\}$ has all moment under $\mathbb{Q}$ for all $e \in \mathbb{Z}^d$ with $|e| = 1$. In particular, γ_e is finite, $\mathbb{Q}$-a.s., thus let $N_x(\omega)$ be $2d$ nearest neighbors of $x \in \mathcal{P}(\omega)$, that is, $\mathcal{N}_x(\omega) := \big\{x + \gamma_e(T_x\omega)e; |e| = 1\big\}$. We call a path $(x_k)_{k=0}^n$ $\mathcal{P}(\omega)$-*nearest neighbor path* if $x_0 \in \mathcal{P}(\omega)$ and $x_k \in \mathcal{N}_{x_{k-1}}$ holds for every $1 \leq k \leq n$.

For each $\omega \in \Omega$, the *simple random walk on the discrete point process* (*RWDPP* for short) is the following Markov chain $((X_n)_{n=0}^\infty, (P_\omega^x)_{x\in\mathcal{P}(\omega)})$ with the state space $\mathcal{P}(\omega)$: for each $z \in \mathcal{P}(\omega)$, let $P_\omega^z(X_0 = z) = 1$ and

$$P_\omega^z(X_{n+1} = y|X_n = x) = \begin{cases} 0 & , y \notin \mathcal{N}_x(\omega), \\ \frac{1}{2d} & , y \in \mathcal{N}_x(\omega). \end{cases} \tag{1.1}$$

We call P_ω^x the *quenched law* and denote by E_ω^x the expectation with respect to P_ω^x.

Our main result is to prove that for $\mathbb{P}$-a.s. ω, the following linear interpolation of $(X_n)_{n=0}^\infty$ converges weakly to a Brownian motion:

$$B_n(t) := \frac{1}{\sqrt{n}}\left(X_{\lfloor tn\rfloor} + (tn - \lfloor tn\rfloor)(X_{\lfloor tn\rfloor+1} - X_{\lfloor tn\rfloor})\right), \qquad t \geq 0. \tag{1.2}$$

Fix $T > 0$ and let $(C[0,T], \mathcal{W}_T)$ be the space of continuous functions $f : [0,T] \to \mathbb{R}$ equipped with the σ-field $\mathcal{W}_T$ of Borel sets relative to the topology introduced by the supremum norm. The precise statement of our main result is as follows:

Theorem 1.1 There is a set $\widehat{\Omega} \subset \Omega_0$ with $\mathbb{P}(\widehat{\Omega}) = 1$ such that on $\widehat{\Omega}$, for all $T > 0$ the law of $(B_n(t))_{0\leq t\leq T}$ on $(C[0,T], \mathcal{W}_T)$ converges, as $n \to \infty$, weakly to the law of a Brownian motion $(B_t)_{0\leq t\leq T}$ with a diffusion matrix D that is independent of ω.

Remark 1.2 In general, we do not find whether the diffusion matrix D is non-degenerate. However, [5, Theorem 4.6] and the proofs of [2, Theorem 6.2] or [3, Theorem 2.1] ensure that the diffusion matrix D is non-degenerate if $\mathbb{P}$ is either reflection invariant or isotropic, that is, invariant with respect to $\mathbb{Z}^d$ a rotations by $\pi/2$.

Let us now describe how the present article is organized. Section 2 recalls the corrector, which was introduced in [2] to establish the quenched invariance principle on percolation clusters. Its properties was analyzed in more detail by [3]. Proposition 2.1 provide the sufficient conditions for the sublinearity of the corrector to prove the quenched invariance principle.

In Section 2 we give the proof of our main result Theorem 1.1. We adopt the approach in [3] and therefore it suffices to prove the sublinearity of the corrector. Therefore, the proof of Theorem 1.1 is the sketch only, and we direct power towards proving the sublinearity. Propositions 3.1, 3.2, 3.4 and 3.6 guarantee that the sufficient conditions for the sublinearity of the corrector are satisfied in our model.

We close this section with some general notation. Let us denote $|x|_\infty := \max_{1\le i\le d}|x_i|$ for $x = (x_1, \dots, x_d) \in \mathbb{Z}^d$ and define $B_\infty(x, n) := \{y \in \mathbb{Z}^d; |x - y|_\infty \le n\}$. The $L^2(\mathbb{P})$-norm for random variables is denoted by $||\cdot||_2$. The canonical unit vectors of $\mathbb{R}^d$ are $e_1, \dots, e_d$.

2 Corrector

In this section, we introduce 'corrector', which plays a key rule in the proof of the quenched invariance principle. Let us first mention the following proposition. For the proof of the proposition we refer the reader to [7, Sections 7 and 9].

Proposition 2.1 There exists a function $\chi : \mathbb{Z}^d \times \Omega_0 \to \mathbb{R}^d$ such that the following properties hold:

(i) (Shift invariance) For $\mathbb{P}$-a.s. ω,

$$\chi(x,\omega) - \chi(y,\omega) = \chi(x-y, T_y\omega), \qquad x,\, y \in \mathcal{P}(\omega).$$

(ii) (Harmonicity) For $\mathbb{P}$-a.s. ω, the function

$$x \longmapsto \chi(x,\omega) + x$$

is harmonic on $\mathcal{P}(\omega)$ with respect to the transition probability (1.1), that is, for every $x \in \mathcal{P}(\omega)$,

$$\frac{1}{2d}\sum_{y\in\mathcal{N}_x(\omega)} (y + \chi(y,\omega)) = x + \chi(x,\omega).$$

(iii) (Square integrability) There exists a positive constant c_1 such that for all $x \in \mathbb{Z}^d$,

$$\sum_{y\in\mathbb{Z}^d} ||(\chi(x,\cdot) - \chi(y,\cdot))\mathbb{1}_{\{x\in\mathcal{P}\}}\mathbb{1}_{\{y\in\mathcal{N}_x\}}||_2^2 \le c_1.$$

(iv) (Sublinearity on average) For every $\epsilon > 0$ and for $\mathbb{P}$-a.s. ω,

$$\lim_{n\to\infty}\frac{1}{n^d}\sum_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \mathbb{1}_{\{|\chi(x,\omega)|\ge\epsilon n\}} = 0.$$

We call the above function χ the *corrector*. It is easy to check from (ii) of Proposition 2.1 that X_n is decomposed in to the difference between a P_ω^0-martingale M_n and $\chi(X_n)$, that is, for $n \ge 0$,

$$M_n^{(\omega)} := X_n + \chi(X_n,\omega). \tag{2.1}$$

For this reason, to show the quenched invariance principle for $(X_n)_{n=0}^\infty$ we adopt the method to use the martingale functional CLT for $(M_n)_{n=0}^\infty$ and an estimate on the corrector χ. In order to get our desired estimate of the corrector χ, it is enough to show its sublinearity, see for the proof of Theorem 1.1 of [2, page 110]:

$$\lim_{n\to\infty} \max_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \frac{|\chi(x,\omega)|}{n} = 0, \qquad \mathbb{P}\text{-a.s.}\,\omega,$$

We now state sufficient condition for the sublinearity of the corrector χ. Let $(N_t)_{t\ge 0}$ be the standard Poisson process with parameter one and we consider the continuous-time Markov chain $(Y_t := X_{N_t})_{t\ge 0}$. Then the following proposition gives sufficient condition for the sublinearity of the corrector. Its proof is given by [8, Theorem 7.15] for i.i.d. configurations, however by [3, Remark 5.5] we can come up with a modification of the proof that covers general ergodic configurations as well.

Proposition 2.2 Suppose that a function $\psi(\cdot,\omega) : \mathcal{P}(\omega) \to \mathbb{R}^d$ satisfies the following conditions (i)-(v) for $\mathbb{P}$-a.s. ω:

(i) (Harmonicity) The function

$$x \longmapsto \psi(x,\omega) + x$$

is harmonic on $\mathcal{P}(\omega)$ with respect to the transition probability (1.1).

(ii) (Sublinearity on average) For every $\epsilon > 0$,

$$\lim_{n\to\infty} \frac{1}{n^d} \sum_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \mathbb{1}_{\{|\psi(x,\omega)|\ge \epsilon n\}} = 0.$$

(iii) (Polynomial growth) There exists a deterministic $\theta > 0$ such that

$$\lim_{n\to\infty} \max_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \frac{|\psi(x,\omega)|}{n^\theta} = 0.$$

(iv) (Diffusive upper bounds) For a deterministic sequence $b_n = o(n^2)$,

$$\sup_{n\ge 1} \max_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \sup_{t\ge b_n} t^{d/2} P_\omega^x(Y_t = x) < \infty. \tag{2.2}$$

$$\sup_{n\ge 1} \max_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \sup_{t\ge b_n} \frac{E_\omega^x[|Y_t - x|]}{\sqrt{t}} < \infty, \tag{2.3}$$

(v) (Control of big jumps) Let $\tau_n := \inf\{t \ge 0; |Y_t - Y_0| \ge n\}$. There exist $c_2 = c_2(\omega) \ge 1$ and $N = N(\omega) \ge 1$ such that for all $t > 0$ and $n \ge N$,

$$\max_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} P_\omega^x(|Y_{t\wedge\tau_n} - x| > c_2 n) = 0.$$

Under these conditions, the function ψ satisfies the sublinearity, that is,

$$\lim_{n\to\infty} \max_{\substack{x\in\mathcal{P}(\omega)\\ |x|\le n}} \frac{|\psi(x,\omega)|}{n} = 0, \qquad \mathbb{P}\text{-a.s.} \tag{2.4}$$

3 Proof of main result

The aim of this section is to prove Theorem 1.1. Let us first give the sketch of the proof, see [2, Sections 6.1 and 6.2] or [3, Theorem 2.1] for more details. We now suppose the following conditions:

- **(S)** The corrector χ satisfies the sublinearity (2.4),
- **(B)** For $\mathbb{P}$-a.s. ω and each $n \geq 0$ there exists a positive constant $K(\omega, n)$ such that $X_n \leq K(\omega, n)$ holds, P_ω^0-a.s.

Let $(M_n^{(\omega)})_{n=0}^\infty$ be a martingale in (2.1) and set $\mathcal{F}_k := \sigma(X_0, \dots, X_k)$. Fix a vector $a \in \mathbb{R}^d$. $(M_n^{(\omega)})_{n=0}^\infty$ is an $L^2(P_\omega^0)$-martingale since the condition (B) implies that X_n and $\chi(X_n, \omega)$ are bounded under P_ω^0. Hence for $\epsilon > 0$ and $m \leq n$, we can define the random variable

$$V_{n,m}^{(\omega)}(\epsilon) := \frac{1}{n} \sum_{k=0}^{m} E_\omega^0 \Big[(a \cdot (M_{k+1}^{(\omega)} - M_k^{(\omega)}))^2 \, \mathbb{1}_{\{|a \cdot (M_{k+1}^{(\omega)} - M_k^{(\omega)})| \geq \epsilon \sqrt{n}\}} \Big| \mathcal{F}_k \Big].$$

Denoting for $K \geq 0$,

$$f_K(\omega) := E_\omega^0 \Big[(a \cdot M_1^{(\omega)})^2 \, \mathbb{1}_{\{|a \cdot M_1^{(\omega)}| \geq K\}} \Big],$$

then from [2, Lemma 6.1] we may write

$$V_{n,m}^{(\omega)}(\epsilon) := \frac{1}{n} \sum_{k=0}^{m} f_{\epsilon\sqrt{n}} \circ T_{X_k}(\omega).$$

Since the Markov chain on environments, $n \to T_{X_n}\omega$ is ergodic, see [2, Theorem 3.2], thus the conditions of the Lindeberg-Feller Martingale CLT hold, see for instance [6, Theorem 7.7.3]. Thereby we conclude that the random continuous function

$$t \mapsto \frac{1}{\sqrt{n}} \Big(a \cdot M_{\lfloor nt \rfloor}^{(\omega)} + (nt - \lfloor nt \rfloor) a \cdot (M_{\lfloor nt \rfloor + 1}^{(\omega)} - M_{\lfloor nt \rfloor}^{(\omega)}) \Big)$$

converges weakly to Brownian motion with mean zero and covariance

$$\mathbb{E}[f_0] = \mathbb{E}[E_\omega^0[(a \cdot M_1^{(\omega)})^2]],$$

which is finite from (iii) of Proposition 2.1. This can be written as $a \cdot Da$ where D is the matrix with coefficients

$$D_{ij} := \mathbb{E}[E_\omega^0[(e_i \cdot M_1^{(\omega)})(e_j \cdot M_1^{(\omega)})]].$$

Applying the Cramér-Wold device, see for instance [6, Theorem 2.9.5], we conclude that the linear interpolation of $t \mapsto M_{\lfloor nt \rfloor}^{(\omega)}/\sqrt{n}$ scales to d-dimensional Brownian motion with covariance matrix D. The condition (S) implies that $M_n^{(\omega)} - X_n = \chi(X_n, \omega) = o(\sqrt{n})$, and so the same conclusion applies to $t \mapsto B_n(t)$ in (1.2).

For the completeness of the proof of Theorem 1.1, we have to show that the conditions (S) and (B) are satisfied. Thanks to (ii) and (iv) of Proposition 2.1, for the proof of (S) it is enough to check the conditions (iii), (iv) and (v) of Proposition 2.2. Let us first prove the conditions (B) and (v) of Proposition 2.2.

Proposition 3.1 The conditions (B) and (v) of Proposition 2.2 hold.

Proof. Let us first construct a set $\widehat{\Omega}_1 \subset \Omega_0$ of full $\mathbb{P}$-measure such that the following holds for $\omega \in \widehat{\Omega}_1$: there is a positive integer $M(\omega)$ such that $\gamma_e(T_x\omega) \leq n$ holds for all $n \geq M(\omega)$ and for all $x \in \mathcal{P}(\omega)$, $e \in \mathbb{Z}^d$ with $|x|_\infty = n$, $|e| = 1$. For $n \geq 1$, we define a subset A_n of Ω_0 as

$$A_n := \{\gamma_e \circ T_x > n \text{ for some } x \in \mathcal{P},\, e \in \mathbb{Z}^d \text{ with } |x|_\infty = n,\, |e| = 1\}.$$

Note that there exists a constant $c_3 = c_3(d)$ such that $\#\{x \in \mathbb{Z}^d; |x|_\infty = n\} \leq c_3 n^{d-1}$ for $n \geq 1$. Chebyshev's inequality then implies that for $n \geq 1$ and $\epsilon > 0$,

$$\begin{aligned}
\mathbb{P}(A_n) &\leq \sum_{|x|_\infty = n} \sum_{|e|=1} \mathbb{P}(x \in \mathcal{P},\, \gamma_e \circ T_x > n) \\
&\leq \sum_{|x|_\infty = n} \sum_{|e|=1} \mathbb{Q}(\Omega_0)^{-1} \mathbb{Q}(\gamma_e > n,\, \omega(0) = 1) \\
&\leq c_3 n^{d-1} \sum_{|e|=1} \mathbb{P}(\gamma_e > n) \\
&\leq \frac{c_3 n^{d-1}}{n^{d+\epsilon}} \sum_{|e|=1} \mathbb{E}[\gamma_e^{d+\epsilon}].
\end{aligned}$$

It follows that the sequence $(\mathbb{P}(A_n))_{n=0}^\infty$ is summable, and then we can construct the desired set $\widehat{\Omega}_1$ from the Borel–Cantelli lemma.

Next we shall show that the condition (v) of Proposition 2.2 is satisfied. Let $\omega \in \widehat{\Omega}_1$ and the positive integer $N(\omega)$ is defied by

$$N(\omega) := M(\omega) + \max\{\gamma_e(T_x\omega); x \in B(0, M(\omega)),\, e \in \mathbb{Z}^d \text{ with } |e| = 1\}.$$

Then the following holds for $\omega \in \widehat{\Omega}_1$: $\mathcal{N}_z(\omega) \subset B_\infty(z, 2n)$ for all $n \geq N(\omega)$ and all $x \in \mathcal{P}(\omega) \cap B_\infty(0, n)$, $y \in \mathcal{P}(\omega) \cap B_\infty(x, n)$ and $z \in \mathcal{P}(\omega) \cap B_\infty(y, n)$. This implies that for $\mathbb{P}$-a.s. ω and all $t > 0$, $n \geq N(\omega)$,

$$\max_{\substack{x \in \mathcal{P}(\omega) \\ |x|_\infty \leq n}} P_\omega^x(|Y_{t \wedge \tau_n} - Y_0|_\infty > 3n) = 0.$$

Replacing $|\cdot|_\infty$ by $|\cdot|$, we find that the condition (v) of Proposition 2.2 is satisfied.

Finally, let us prove the condition (B). For fixed $\omega \in \widehat{\Omega}_1$ and $n \geq 0$ we consider a $\mathcal{P}(\omega)$-nearest neighbor path $(0 = x_0, x_1, \ldots, x_n)$ starting at the origin of length n. It is clear by the choice of $N(\omega)$ that $|x_1|_\infty \leq N(\omega)$ holds. Thanks to the choice of $N(\omega)$, $|x_2|_\infty$ has to be less than $2N(\omega)$. By induction on the steps of $(0 = x_0, x_1, \ldots, x_n)$ we can see that for $2 \leq i \leq n$, $|x_i| \leq 2^{i-2}N(\omega)$. Hence we get the following rough upper bound on the $|\cdot|_\infty$-distance of x_n:

$$|x_n|_\infty \leq N(\omega)\left(1 + \sum_{i=2}^n 2^{i-2}\right) = 2^{n-1}N(\omega),$$

which proves the condition (B). □

We next prove the bound (2.2) in the condition (iv) of Proposition 2.2.

Proposition 3.2 The bound (2.2) in the condition (iv) of Proposition 2.2 holds.

Proof. From [7, Claim 5.9] there exist positive constants c_3' and c_4' depending only on d such that the following holds, $\mathbb{P}$-a.s.: a positive integer N such that for sufficiently large n and all x, $y \in \mathcal{P}$,

$$P_\omega^x(X_n = y) \le \frac{c_3'}{(n - c_4')^{d/2}}.$$

By standard arguments we get the corresponding bound for the discrete time version of $(Y_t)_{t\ge 0}$. This proves the theorem. □

For (2.3) in the condition (iv) of Proposition 2.2, let us prepare some notation and lemma. We denote by $d_\omega(x, y)$ the graph distance on $\mathcal{P}(\omega)$. The random walk $|\cdot|$ $(Y_t)_{t\ge 0}$ has non-bounded jumps under the Euclidean distance, however it has bounded jumps under the graph distance d_ω, so the same strategy as in [1, Proposition 3.4] or [3, Proposition 6.2] that we get

$$\sup_{z\in\mathbb{Z}^d} \sup_{t\ge 1} \frac{E_\omega^z[d_\omega(z, Y_t)]}{\sqrt{t}} < \infty. \tag{3.1}$$

For this reason, if $d_\omega(x, y)$ is comparable with the Euclidean distance $|x - y|$, then we can get the corresponding bound for the Euclidean distance version.

Lemma 3.3 There exist positive constants ρ, c_5, and c_6 depending only on d such that

$$\mathbb{Q}(0, x \in \mathcal{P},\ d_\omega(0, x) \le \rho|x|) \le c_5 \exp\{-c_6|x|\}.$$

Proof. We consider the lattice cubes $B_L(x) := x + [-L, L)^d$, $L \in \mathbb{N}$, $x \in \mathbb{Z}^d$. We call $B_L(x)$ *blocked* if every $\mathcal{P}$-nearest neighbor path crossing $B_L(x)$ does not pass through $B_L(x)$, that is, for all $1 \le i \le d$, $j \in [1, d] \setminus \{i\}$ and $-L \le k_j < L$, there exists $-L \le m_i < L$ such that

$$\omega(x + k_1e_1 + \cdots + m_ie_i + \cdots + k_de_d) = 1.$$

If $B_L(x)$ is not blocked, then we call it *unblocked*. From the assumptions (A1)-(A3), we can see that $p_L := \mathbb{Q}(B_L(0)$ is unblocked) converges to zero as $L \to \infty$. For $L \in \mathbb{N}$ we set $\mathcal{C}_L(0) := \{B_L(x); x \in 4L\mathbb{Z}^d\}$ and then let us introduce for $1 \le k \le d$ and distinct unit vectors $v_1, \dots, v_k \in \{e_1, \dots, e_d\}$,

$$\mathcal{C}_L(v_1, \dots, v_k) := \{B_L(x); x \in 2L(v_1 + \cdots + v_k) + 4L\mathbb{Z}^d\}.$$

To simplify notation, let $\mathcal{C}_L^{(i)}$ $(1 \le i \le 2^d)$ be an enumeration of $\mathcal{C}_L(0)$ and $\mathcal{C}_L(v_1, \dots, v_k)$'s. In addition, we define $\mathcal{I}_L := \bigcup_{i=1}^{2^d} \mathcal{C}_L^{(i)}$. Choose $\delta := 1/2^d$ and let us fix $\ell \le L \in \mathbb{N}$ and $n \ge 0$. For a lattice animal $\mathcal{A}_n$ on $2L\mathbb{Z}^d$ of size n and containing 0, the event $\Gamma(x, \mathcal{A}_n)$ is defined as follows: $d_\omega(0, x) < \delta|x|_\infty/(2L)$ and a $\mathcal{P}(\omega)$-nearest neighbor path realizing $d_\omega(0, x)$, which is chosen by a deterministic algorithm, intersects with the n elements of $\mathcal{I}_L$ and is included in $\bigcup_{y\in\mathcal{A}_n} B_L(x)$. Furthermore, let $G(k, i, \mathcal{A}_n)$ be the event that $\bigcup_{y\in\mathcal{A}_n} B_L(y)$ contains the k elements of $\mathcal{C}_L^{(i)}$ and at least δn of its k elements of $\mathcal{C}_L^{(i)}$ are



Figure 1: A $\mathcal{P}(\omega)$-nearest neighbor path realizing $d_\omega(0,x)$ and a lattice animal $\mathcal{A}_n$. The crosses stand for sites on a $\mathcal{P}(\omega)$-nearest neighbor path.

unblocked. Noting that $\mathcal{P}(\omega)$-nearest neighbor paths realizing $d_\omega(0,x)$ intersect at least $|x|_\infty/(2L)$ elements of $\mathcal{I}_L$, we have

$$\Gamma(x,\mathcal{A}_n) \subset \bigcup_{n/2^d \le k \le n} \bigcup_{i=1}^{2^d} G(k,i,\mathcal{A}_n).$$

On the other hand, for each $1 \le i \le 2^d$, the events $\{C \text{ is unblocked}\}$ for $C \in \mathcal{C}_L^{(i)} \subset \bigcup_{y\in\mathcal{A}_n} B_L(x)$ are independent and have the same probability p_L. Set $\delta'(L) := \delta - p_L$ and

$$\alpha(L) := (p_L + \delta'(L)) \log \frac{p_L + \delta'(L)}{p_L} + (1 - p_L - \delta'(L)) \log \frac{1 - p_L - \delta'(L)}{1 - p_L}.$$

The Chernoff bound yields

$$\mathbb{Q}(G(k,i,\mathcal{A}_n)) \le \exp\{-\alpha(L)k\}.$$

Hence we have

$$\begin{aligned} \mathbb{Q}(\{0, x \in \mathcal{P}\} \cap \Gamma(x,\mathcal{A}_n)) &\le \sum_{n/2^d \le k \le n} \sum_{i=1}^{2^d} \mathbb{Q}(G(k,i,\mathcal{A}_n)) \\ &\le 2^d n \exp\{-\alpha(L)\delta n\}. \end{aligned}$$

Since $\alpha(L) \to \infty$ as $L \to \infty$, we can find L such that $(2d)^{2n} 2^d n \exp\{-\alpha(L)\delta n\} \le 2^d n e^{-2n}$ for $n \ge 1$. Recall that the number of lattice animals on $2L\mathbb{Z}^d$, of size n, containing the origin, is roughly bounded from above by $(2d)^{2n}$, see [4, Lemma 1]. It follows that we get

$$\mathbb{Q}(0, x \in \mathcal{P},\, d_\omega(0,x) < \delta|x|_\infty/(2L))$$

$$\begin{aligned}
&\le \sum_{n \ge |x|_\infty/(2L)} (2^d)^{2n} 2^d n \exp\{-\alpha(L)\delta n\} \\
&\le 2^d \sum_{n \ge |x|_\infty/(2L)} \exp\{-n\} \\
&\le \frac{2^d}{1-e^{-1/2}} \exp\{-|x|_\infty/(2L)\},
\end{aligned}$$

which shows the conclusion. □

Under the above preparation, let us show (2.3) in (iv) of Proposition 2.2.

Proposition 3.4 The bound (2.3) in (iv) of Proposition 2.2 holds.

Proof. We first show that there are a positive constant c_7 depending only on d and $\widehat{\Omega}_2 \subset \Omega$ of full $\mathbb{Q}$-measure such that the following holds for $\omega \in \widehat{\Omega}_2$: there exists a positive integer $N(\omega)$ such that $d_\omega(z,y) \ge \rho|z-y|$ holds for all $n \ge N(\omega)$ and all $z, y \in \mathcal{P}(\omega)$ with $|z| \le n$, $|z-y| \ge c_7 \log n$. Set $c_7 := 2(d+2)/c_6$. From Lemma 3.3, we have for some positive constant c_8 depending only on d and for all N sufficiently large,

$$\begin{aligned}
&\sum_{n=N}^{\infty} \sum_{\substack{z,y \in \mathbb{Z}^d \\ |z| \le n \\ |z-y| \ge c_7 \log n}} \mathbb{Q}(0, y-z \in \mathcal{P},\, d_\omega(0,y-z) < \rho|z-y|) \\
&\le \sum_{n=N}^{\infty} \sum_{\substack{z,y \in \mathbb{Z}^d \\ |z| \le n \\ |z-y| \ge c_7 \log n}} c_5 \exp\{-c_6|z-y|\} \\
&\le \sum_{n=N}^{\infty} \sum_{|z| \le n} \sum_{K \ge c_7 \log n} c_8 K^{d-1} \exp\{-c_6 K\}.
\end{aligned}$$

It is easy to see that this sum converges. By the Borel–Cantelli lemma the assertion stated in the beginning of the proof is verified.

Fix $\omega \in \widehat{\Omega}_2$. Then we obtain for all $n \ge N(\omega)$, $|z| \le n$ and $t \ge n$,

$$\begin{aligned}
\frac{E_\omega^z[|z-Y_t|]}{\sqrt{t}} &\le \frac{1}{\sqrt{t}} \bigl(\rho^{-1} E_\omega^z[d_\omega(z,Y_t)] + c_7 \log n\bigr) \\
&\le \rho^{-1} \frac{E_\omega^z[d_\omega(z,Y_t)]}{\sqrt{t}} + c_7 \frac{\log n}{n},
\end{aligned}$$

which shows the bound (2.3) in (iv) of Proposition 2.2. □

Finally, we will prove that the corrector χ satisfies Proposition 2.2 (iii). To do this, we prepare some notation and lemma. Let us denote by $E_{\theta,n}$ the event that for any $y \in \mathcal{P} \cap [-n,n]^d$, there exists a $\mathcal{P}$-nearest neighbor path $(0 = z_0, z_1, \dots, z_m = y)$ from 0 to y such that $\max_{0 \le k \le m} |z_k|_\infty \le n^\theta$.

Lemma 3.5 Suppose that the following condition (C) holds:

(C) For some $\lambda > 0$ the sequence $(\mathbb{P}(E_{\theta,n}^c))_{n=1}^{\infty}$ is summable.

Then the condition (iii) of Proposition 2.2 holds.

Proof. For $\omega \in \Omega_0$ let

$$R_n(\omega) := \max_{\substack{x \in \mathcal{P}(\omega) \\ |x|_\infty \le n}} |\chi(x, \omega)|.$$

From Proposition 2.1 (iii) we have

$$\begin{aligned}
&\mathbb{E}[R_n^2 \mathbb{1}_{E_{\theta,n}}] \\
&\le (4\lambda n)^2 \sum_{|x|_\infty \le n^\theta} \sum_{y \in \mathbb{Z}^d} \mathbb{E}[|\chi(x+y, \cdot) - \chi(x, \cdot)|^2 \mathbb{1}_{\{x \in \mathcal{P}\}} \mathbb{1}_{\{y \in \mathcal{N}_x\}}] \\
&\le c_9 n^{\theta+2}
\end{aligned}$$

for every $n \ge 1$ and some constant $c_9 = c_9(\lambda)$. Applying Chebyshev's inequality, we obtain for $\theta' > 0$,

$$\mathbb{P}(R_n \mathbb{1}_{E_{\theta,n}} \ge n^{\theta'}) \le c_9 n^{-2\theta' + \theta + 2}.$$

Therefore the condition (C) yields that for $\theta' > (\theta + 3)/2$,

$$\begin{aligned}
\sum_{n=0}^\infty \mathbb{P}(R_n \ge n^{\theta'}) &\le \sum_{n=0}^\infty \mathbb{P}(R_n \mathbb{1}_{E_{\theta,n}} \ge n^{\theta'}) + \sum_{n=0}^\infty \mathbb{P}(E_{\theta,n}^c) \\
&\le 1 + c_9 \sum_{n=1}^\infty n^{-2\theta'+\theta+2} + \sum_{n=0}^\infty \mathbb{P}(E_{\theta,n}^c) < \infty,
\end{aligned}$$

which proves the condition (iii) of Proposition 2.2 by the Borel–Cantelli lemma. □

Thanks to the above lemma, for the proof of the condition (iii) of Proposition 2.2 it suffices to check the condition (C). Let us start to prove the condition (C). For $x = (x_1, \dots, x_d) \in \mathbb{Z}^d$ and $2 \le i \le d$, let us denote by $F_{i,n}(x)$ the event that for any $y \in \mathcal{P} \cap [-n, n]^i \times \{(x_{i+1}, \dots, x_d)\}$, there exists a $\mathcal{P}$-nearest neighbor self-avoiding path $(x = r_0, r_1, \dots, r_m = y)$ from x to y such that $r_k \in [-n, n]^i \times \{(x_{i+1}, \dots, x_d)\}$, $1 \le k \le m$. In addition, the event $G_{i,n}(x)$ is defined by the event that the set $[-n, n]^i \times \{(x_{i+1}, \dots, x_d)\}$ appearing in the statement of the definition of $F_{i,n}(x)$ is replaced by $\{x_1\} \times [-n, n]^i \times \{(x_{i+2}, \dots, x_d)\}$.

Proposition 3.6 The condition (C) is satisfied.

Proof. Let $\rho := 1 - \mathbb{Q}(\Omega_0)$. Suppose that for some $2 \le i \le d - 1$ there exist positive constants $C_1^{(i)}, C_2^{(i)}$ such that for n large enough and all $x = x_1 e_1 + \dots + x_i e_i \in \mathbb{Z}^d \cap [-n, n]^d$,

$$\mathbb{Q}(F_{i,n}(x)^c \cap \Omega_x) + \mathbb{Q}(G_{i,n}(x)^c \cap \Omega_x) \le C_1^{(i)} \rho^{C_2^{(i)} n}. \tag{3.2}$$

Denote by $H_{i,n}(z)$ the event that for all $-n \le k \le n$ there is a site $w = k e_1 + w_2 e_2 + \dots + w_i e_i + z_{i+1} e_{i+1}$ such that $w \in \mathcal{P}$ and $-n \le w_j \le n$ for $2 \le j \le i$. A straightforward calculation then shows that we get for $z = z_1 e_1 + \dots + z_{i+1} e_{i+1} \in \mathbb{Z}^d \cap [-n, n]^d$,

$$\begin{aligned}
\mathbb{Q}(F_{i+1,n}(z)^c \cap \Omega_z) &\le \mathbb{Q}(F_{i,n}(z)^c \cap \Omega_z) + \mathbb{Q}(H_{i,n}(z)^c) \\
&\quad + \mathbb{Q}(F_{i+1,n}(z)^c \cap \Omega_z \cap F_{i,n}(z) \cap H_{i,n}(z)).
\end{aligned} \tag{3.3}$$

By (3.2) the first term of the right-hand side of (3.3) is less than $C_1^{(i)}\rho^{C_2^{(i)}n}$. Moreover, it is easy to see that there are positive constants c_{10} and c_{11} such that for n sufficiently large, the second term of the right-hand side of (3.3) is bounded from above by $c_{10}\rho^{c_{11}n}$. Let us estimate the third term of the right-hand side of (3.3). On the event $H_{i,n}(z)$, there are sites $u_k \in \mathcal{P}$, $-n \le k \le n$ such that

$$u_k \in \mathcal{U}(i,n,k) := \{ke_1 + v_2e_2 + \cdots + v_ie_i + z_{i+1}e_{i+1}; -n \le v_2, \dots, v_i \le n\}.$$

Note that on the event

$$I_{i,n}(z) := \Omega_z \cap F_{i,n}(z) \cap H_{i,n}(z) \cap \bigcap_{k=-n}^{n} (G_{i,n}(u_k) \cap \Omega_{u_k}),$$

for each $y = y_1e_1 + \cdots + y_{i+1}e_{i+1} \in [-n,n]^d \cap \mathcal{P}$ the site u_{y_1} is connected to y by a $\mathcal{P}$-nearest neighbor path included in $\{y_1\} \times [-n,n]^i \times \{0\}^{d-i-1}$ and is connected to z by a $\mathcal{P}$-nearest neighbor path included in $[-n,n]^i \times \{0\}^{d-i}$. For this reason, the event $F_{i+1,n}(z)$ occurs on the event $I_{i,n}(z)$. We hence obtain

$$\begin{aligned}
&\mathbb{Q}(F_{i+1,n}(z)^c \cap \Omega_z \cap F_{i,n}(z) \cap H_{i,n}(z)) \\
&\le \sum_{\substack{(u_{-n},\dots,u_n) \\ \in \mathcal{U}(i,n,-n)\times\cdots\times\mathcal{U}(i,n,n)}} \mathbb{Q}\left(F_{i+1,n}(z)^c \cap \Omega_z \cap F_{i,n}(z) \cap H_{i,n}(z) \cap \bigcap_{k=-n}^{n} \Omega_{u_k} \right) \\
&\le \sum_{\substack{(u_{-n},\dots,u_n) \\ \in \mathcal{U}(i,n,-n)\times\cdots\times\mathcal{U}(i,n,n)}} \left(\mathbb{Q}(F_{i+1,n}(z)^c \cap I_{i,n}(z)) + \sum_{j=-n}^{n} \mathbb{Q}(G_{i,n}(u_j)^c \cap \Omega_{u_j}) \right) \\
&\le C_1^{(i)}(2n+1)^{i+1}\rho^{C_2^{(i)}n},
\end{aligned}$$

and therefore the following holds:

$$\mathbb{Q}(F_{i+1,n}(z)^c \cap \Omega_z) \le C_1^{(i)}\rho^{C_2^{(i)}n} + c_{10}\rho^{c_{11}n} + C_1^{(i)}(2n+1)^{i+1}\rho^{C_2^{(i)}n}.$$

By the same argument as the above, we can estimate $\mathbb{Q}(G_{i+1,n}(z)^c \cap \Omega_z)$, so there are some positive constants $C_1^{(i+1)}$ and $C_2^{(i+1)}$ such that for n large enough and all $z = z_1e_1 + \cdots + z_{i+1}e_{i+1} \in \mathbb{Z}^d \cap [-n,n]^d$,

$$\mathbb{Q}(F_{i+1,n}(z)^c \cap \Omega_z) + \mathbb{Q}(G_{i+1,n}(z)^c \cap \Omega_z) \le C_1^{(i+1)}\rho^{C_2^{(i+1)}n}.$$

By induction on i it is enough to show (3.2) in the case $i = 2$. To simplify notation, let $d = 2$. It suffices to get the bound for $\mathbb{Q}(F_{2,n}(x)^c \cap \Omega_x)$, $x = x_1e_1 + x_2e_2 \in \mathbb{Z}^2 \cap [-n,n]^2$, since an analogous statement with $\mathbb{Q}(G_{2,n}(x)^c \cap \Omega_x)$ instead of $\mathbb{Q}(F_{2,n}(x)^c \cap \Omega_x)$ can be derived similarly. Without loss of generality we can assume $x_2 \ge 0$. To this end, for $n \ge 1$, $u \in [0,n] \cap \mathbb{Z}$, $0 < \delta < 1$ and $L \ge 1$ let us introduce the events

$$\begin{aligned}
\Lambda_0(u,L) &:= \{(u-j)e_2 \in \mathcal{P} \text{ for some } 1 \le j \le L\}, \\
\Lambda_1(u,\delta,L,n) &:= \left\{ \frac{1}{2n+1} \sum_{i=-n}^{n} \mathbb{1}_{\Lambda_0(u,L)} \circ T_{ie_1} > 1 - \delta \right\},
\end{aligned}$$

and $\Lambda_2(u, L, n)$ defined as follows: for all $\ell \le j \le L$, there is an $0 \le i_j \le n$ such that

$$\omega(i_j e_1) = \omega(i_j e_1 + (u - j)e_2) = 1,$$

and for all $0 \le m \le \ell - 1$, there is a $0 \le k_m \le n$ such that

$$\omega(k_m e_1 + (u - m)e_2) = \omega(k_m e_1 + (u - (m + \ell))e_2) = 1.$$

The event $\Lambda_1(u, \delta, L, n)$ means that almost all vertical lines are blocked in the slab $[-n, n] \times [u - L, u]$ and the event $\Lambda_2(u, L, n)$ means that all lines in the slab $[-n, n] \times [u - L, u]$ are connected by $\mathcal{P}$-nearest neighbor paths included in $[-n, n]^2$. We will estimate $\mathbb{Q}(\Lambda_1(u, \delta, L, n)^c)$. Let $0 < \delta < \mathbb{Q}(\Omega_0)$ and we choose $0 < \delta_1 < \delta$. Observe that if

$$n > \frac{1}{2}\left(\frac{(1 - \delta_1)\ell}{\delta - \delta_1} - 1\right), \tag{3.4}$$

then

$$\frac{2n + 1}{\lfloor (2n + 1)/\ell \rfloor} \, \frac{1 - \delta}{\ell} < 1 - \delta_1.$$

Let $I_k := \{-n + k + m\ell; 0 \le m \le \lfloor (2n + 1)/\ell \rfloor - 1\}$, $0 \le k \le \ell - 1$. We get for any n satisfying (3.4),

$$\begin{aligned}
&\mathbb{Q}(\Lambda_1(u, \delta, L, n)^c) \\
&\le \mathbb{Q}\left(\frac{1}{2n + 1} \sum_{k=0}^{\ell-1} \sum_{i \in I_k} \mathbb{1}_{\Lambda_0(u,L)} \circ T_{ie_1} \le 1 - \delta\right) \\
&\le \sum_{k=0}^{\ell-1} \mathbb{Q}\left(\frac{1}{2n + 1} \sum_{i \in I_k} \mathbb{1}_{\Lambda_0(u,L)} \circ T_{ie_1} \le \frac{1 - \delta}{\ell}\right) \\
&\le \ell \, \mathbb{Q}\left(\frac{1}{\lfloor (2n + 1)/\ell \rfloor} \sum_{i \in I_0} \mathbb{1}_{\Lambda_0(x,L)} \circ T_{ie_1} \le \frac{2n + 1}{\lfloor (2n + 1)/\ell \rfloor} \, \frac{1 - \delta}{\ell}\right) \\
&\le \ell \, \mathbb{Q}\left(\frac{1}{\lfloor (2n + 1)/\ell \rfloor} \sum_{i \in I_0} \mathbb{1}_{\Lambda_0(u,L)} \circ T_{ie_1} < 1 - \delta_1\right).
\end{aligned}$$

It is clear from (A2) and (A3) that we can take numbers $0 < \delta_2 < 1$ and $L \ge 2\ell - 1$ with $1 - \delta_1 \le (1 - \delta_2) \, \mathbb{Q}(\Lambda_0(0, L))$. Thus the most right-hand side of the above expression is less than

$$\ell \, \mathbb{Q}\left(\frac{1}{\lfloor (2n + 1)/\ell \rfloor} \sum_{i \in I_0} \mathbb{1}_{\Lambda_0(u,L)} \circ T_{ie_1} < (1 - \delta_2) \, \mathbb{Q}(\Lambda_0(0, L))\right). \tag{3.5}$$

Using the Chernoff bound, we estimate (3.5) from above by

$$\ell \exp\left\{-\frac{\delta_2^2 \, \mathbb{Q}(\Lambda_0(0, L))}{2} \left\lfloor \frac{2n + 1}{\ell} \right\rfloor\right\}.$$

It follows that

$$\mathbb{Q}(\Lambda_1(u,\delta,L,n)^c) \le \ell \exp\left\{-\frac{\delta_2^2\,\mathbb{Q}(\Lambda_0(0,L))}{2}\left\lfloor\frac{2n+1}{\ell}\right\rfloor\right\}. \tag{3.6}$$

holds for all n satisfying (3.4).

Let us next estimate $\mathbb{Q}(\Lambda_2(x,L,n)^c)$. Noting that by the assumption (A2) $\omega(k\ell e_1)$, $k \ge 0$ are independent under $\mathbb{Q}$, we have for $n \ge \ell$,

$$\begin{aligned}
&\mathbb{Q}(\Lambda_2(u,L,n)^c)\\
&\le \sum_{j=\ell}^{L}\mathbb{Q}\Big(\text{for every } 0 \le i \le n/\ell,\\
&\qquad\qquad \omega(i\ell e_1) = \omega(i\ell e_1 + (u-j)e_2) = 1 \text{ fails}\Big)\\
&\quad + \sum_{m=0}^{\ell-1}\mathbb{Q}\Big(\text{for every } 0 \le k \le n/\ell,\ \omega(k\ell e_1 + (u-m)e_2)\\
&\qquad\qquad = \omega(k\ell e_1 + (u-(m+\ell))e_2) = 1 \text{ fails}\Big)\\
&\le (L-\ell+1)(1-\mathbb{Q}(\Omega_0)^2)^{n/\ell} + \ell(1-\mathbb{Q}(\Omega_0)^2)^{n/\ell}\\
&= (L+1)(1-\mathbb{Q}(\Omega_0)^2)^{n/\ell}.
\end{aligned} \tag{3.7}$$

Let us estimate the left-hand side of (3.2). (3.6) and (3.7) ensure that

$$\begin{aligned}
&\mathbb{Q}(E_{2,n}(x)^c \cap \Omega_x)\\
&\le \mathbb{Q}(\Lambda_1(x_2,\delta,L,n)^c) + \mathbb{Q}(\Lambda_2(x_2,L,n)^c)\\
&\quad + \mathbb{Q}(E_{2,n}(x)^c \cap \Omega_x \cap \Lambda_1(x_2,\delta,L,n) \cap \Lambda_2(x_2,L,n))\\
&\le \exp\left\{-\frac{\delta_2^2\,\mathbb{Q}(\Lambda_0(L))}{2}\left\lfloor\frac{n+1}{\ell}\right\rfloor\right\} + L(1-\mathbb{Q}(\Omega_0)^2)^{n/\ell}\\
&\quad + \mathbb{Q}(E_{2,n}(x)^c \cap \Omega_x \cap \Lambda_1(x_2,\delta,L,n) \cap \Lambda_2(x_2,L,n)).
\end{aligned} \tag{3.8}$$

holds for $x = x_1e_1 + x_2e_2 \in \mathbb{Z}^2 \cap [-n,n]^2$. We now prove that there is a $y_1e_1 + y_2e_2 \in \mathcal{P} \cap [-n,n]^2$ such that

$$\sum_{i=-n}^{n} \mathbb{1}_{\{\omega(ie_1+y_2e_2)=1\}} < \delta(2n+1) \tag{3.9}$$

holds on the event $E_{2,n}(x)^c \cap \Omega_x \cap \Lambda_1(x_2,\delta,L,n) \cap \Lambda_2(x_2,L,n)$. We suppose that on the event $\Omega_x \cap \Lambda_1(x_2,\delta,L,n) \cap \Lambda_2(x_2,L,n)$, (3.9) fails for all $y_1e_1+y_2e_2 \in \mathcal{P} \cap [-n,n]^2$. On the event $\Omega_x \cap \Lambda_1(x_2,\delta,L,n)$, there exist at least $\lceil(1-\delta)(2n+1)^2\rceil$ vertical lines contained in $[-n,n]^2$ such that each of these vertical lines has a site in $\mathcal{P}$ standing for the slab $[-n,n] \times [x_2-L,x_2]$. Moreover, On the event $\Omega_x \cap \Lambda_2(x_2,L,n)$, all lines in the slab $[-n,n] \times [x_2-L,x_2]$ are connected to x by $\mathcal{P}$-nearest neighbor paths included in $[-n,n]^2$. It follows that on the event $\Omega_x \cap \Lambda_1(x_2,\delta,L,n) \cap \Lambda_2(x_2,L,n)$, at least $\lceil(1-\delta)(2n+1)^2\rceil$ vertical lines contained in $[-n,n]^2$ are connected to x by $\mathcal{P}$-nearest neighbor paths. On the other hand, on the event $\Omega_x \cap \Lambda_1(x_2,\delta,L,n) \cap \Lambda_2(x_2,L,n)$ there is a horizontal line contained in $[-n,n]^2$ such that it does not have sites connected to x by a



Figure 2: The event $E_{2,n}(x)^c \cap \Omega_x \cap \Lambda_1(x_2, \delta, L, n) \cap \Lambda_2(x_2, L, n)$. The crosses standing for L horizontal lines are connected to x by $P(\omega)$-nearest neighbor paths included in $[-n, n]^2$. There exists some site $y \in \mathcal{P}(\omega) \cap [-n, n]^2$ such that the site y is not connected to x and solid vertical lines cannot have points in the slab $[-n, n] \times [x_2 - L, x_2]$.

$\mathcal{P}$-nearest neighbor path included in $[-n, n]^2$ and this horizontal line has at least $\lceil \delta(2n+1) \rceil$ sites in $\mathcal{P}$. This means that the number of vertical lines contained in the box $[-n, n]^2$ has to be strictly grater than $(1-\delta)(2n+1)+\delta(2n+1) = 2n+1$, which contradicts that the number of vertical lines contained in the box $[-n, n]^2$ is equal to $2n + 1$. Therefore, on the event $\Omega_x \cap \Lambda_1(x_2, \delta, L, n) \cap \Lambda_2(x_2, L, n)$, (3.9) holds for some $y_1e_1 + y_2e_2 \in \mathcal{P} \cap [-n, n]^2$.

The same argument as in the proof of (3.6) implies

$$\begin{aligned} &\mathbb{Q}(E_{2,n}(x)^c \cap \Omega_x \cap \Lambda_1(x_2, \delta, L, n) \cap \Lambda_2(x_2, L, n)) \\ &\leq (2n+1)^2 \mathbb{Q}\bigg(\sum_{i=-n}^{n} \mathbb{1}_{\{\omega(ie_1)=1\}} < \delta(2n+1) \bigg) \\ &\leq \ell(2n+1)^2 \exp\bigg\{ -\frac{\delta_3^2 \, \mathbb{Q}(\Omega_0)}{2} \left\lfloor \frac{2n+1}{\ell} \right\rfloor \bigg\} \end{aligned} \tag{3.10}$$

for n sufficient large and some $0 < \delta_3 < 1$. According to (3.8) and (3.10), the bound (3.2) holds for $i = 2$. □

Acknowledgments. The author thanks Prof. Shigenori Matsumoto and Prof. Takao Nishikawa for many helpful suggestions. I am grad to thank Prof. Jun Misumi and Prof. Ryoki Fukushima and for discussions and important comments on this problem.

References

[1] Martin T. Barlow. Random walks on supercritical percolation clusters. *Ann. Probab.*, Vol. 32, No. 4, pp. 3024–3084, 2004.

[2] Noam Berger and Marek Biskup. Quenched invariance principle for simple random walk on percolation clusters. *Probab. Theory Related Fields*, Vol. 137, No. 1-2, pp. 83–120, 2007.

[3] Marek Biskup and Timothy M. Prescott. Functional CLT for random walk among bounded random conductances. *Electron. J. Probab.*, Vol. 12, pp. no. 49, 1323–1348, 2007.

[4] J. Theodore Cox, Alberto Gandolfi, Philip S. Griffin, and Harry Kesten. Greedy lattice animals. I. Upper bounds. *Ann. Appl. Probab.*, Vol. 3, No. 4, pp. 1151–1169, 1993.

[5] A. De Masi, P. A. Ferrari, S. Goldstein, and W. D. Wick. An invariance principle for reversible Markov processes. Applications to random motions in random environments. *J. Statist. Phys.*, Vol. 55, No. 3-4, pp. 787–855, 1989.

[6] Richard Durrett. *Probability: theory and examples*. Duxbury Press, Belmont, CA, third edition, 1996.

[7] Ron Rosenthal. Random walk on discrete point processes. *Preprint*, 2010. `http://arxiv.org/PS_cache/arxiv/pdf/1005/1005.1398v1.pdf`.

[8] Kumagai Takashi. Random walks on disordered media and their scaling limits. *Notes on Saint Flour Lectures 2010, preprint*, 2010. `http://www.kurims.kyoto-u.ac.jp/~kumagai/StFlour-TK.pdf`.